

## Supplementary material

### Supplementary Methods

#### ***Adenovirus production***

The Fc domain, containing mutations in the IgG-R and complement binding domains, was amplified by PCR from an IL-10-Fc fusion plasmid (generously provided by Dr Terry Storm, Harvard University, USA). The extracellular domain of mouse N-cadherin was cloned with the Fc domain into pDC515 (Microbix). The secretion signal was cloned from full length N-cadherin onto the Fc fragment and cloned into pDC515. Both were recombined with the adenovirus genomic plasmid by co-transfection into 293 cells. The resultant adenoviruses were termed RAd SNC-Fc and RAd Fc (control). This chimeric molecule comprised of SNC and an antibody constant domain, enables protein A binding, as well as extending plasma half-life<sup>1-3</sup> and possibly increasing receptor–ligand interaction. Such immunoadhesins have use *in vivo* as potential therapeutic agents<sup>4</sup>.

#### ***Purification and culture of mouse blood monocytes***

Mouse peripheral blood monocytes were purified by Ficoll-Hypaque gradient (Ficoll-Paque Plus: Amersham Biosciences), followed by differential adherence and culture in 20ng/ml M-CSF for 7-10 days to induce differentiation into macrophages.

#### ***Purification of rabbit foam cell-macrophages***

New Zealand White rabbits (Harlan, UK) fed a 1% cholesterol-supplemented diet had sterile sponges placed under the dorsal skin to generate foam cell macrophages, as described<sup>5</sup>.

#### ***siRNA Knockdown***

Two HP validated silencing RNA oligonucleotides (siRNA) for FGF-R1 and control (Allstars Negative control siRNA) were purchased from Qiagen (catalogue numbers SI02224677, SI02224684 and 1027281). VSMCs ( $8 \times 10^5$ ) were subjected to Amaxa nucleofection with

250pmol of FGF-R1 or control siRNAs using the VSMC kit and U-25 program following the manufacturer's instructions (Amaya, Inc., Cologne Germany). Treated cells were analysed 24 hours after nucleofection. Knockdown of FGF-R was estimated as  $85\pm 7\%$  ( $n=3$ ,  $p<0.05$ ) by quantitative PCR and was confirmed by immunocytochemistry (data not shown).

### ***Western blotting***

SDS lysed cell extracts were subjected to Western blotting as described previously<sup>6</sup>. Blots were incubated overnight at 4°C with primary antibodies diluted in Starting block (Pierce, Chester, UK). Antibodies were used at the following concentrations: total and pAkt (Cell Signalling, 1:1000). Bound antibodies were detected by rabbit anti-mouse horseradish peroxidase conjugated antibodies (Dako, High Wycombe, UK) and enhanced chemiluminescence (Amersham International, Little Chalfont, UK).

### ***Galactolight assay***

TOPgal transgenic mouse VSMCs were grown from aortic explants, as described previously<sup>7</sup>. These VSMCs contain the  $\beta$ -galactosidase gene under the control of the  $\beta$ -catenin promoter. TOPgal VSMCs were treated with 0.01% (w/v) trypsin, 1 mM  $\text{CaCl}_2$  in PBS) and seeded into a 24 well plate in the presence of SNC-Fc or Fc and Fas-L.  $\beta$ -galactosidase activity was quantified using a chemiluminescent assay called Galactolight as described by the manufacturer's instructions (Tropix).

### ***Aggregation and adhesion assays***

Cell aggregation and cell-cell adhesion were determined as previously<sup>8, 9</sup>.

### ***Invasion of monocytes and proliferation of macrophages***

Mouse peripheral blood monocyte invasion and macrophage proliferation were assessed as previously<sup>10</sup>.

### **Quantitative PCR**

Total RNA was isolated by the RNAeasy kit (Qiagen) was reverse transcribed and subjected to quantitative PCR for N-cadherin and FGF-R using Quantitect primers (Qiagen, QT00102837 and QT00198548, respectively) as described by the manufacturer.

### **In vivo experiments**

#### *Quantification of plasma SNC-Fc and lipoprotein levels*

Plasma samples were taken at 2, 6, 8, 14 and 28 days after RAd administration and levels of SNC-Fc were analysed by ELISA as described previously<sup>6</sup>. Plasma lipid profiles were analyzed in terminal plasma samples as previously described<sup>11</sup>.

#### *Immunohistochemistry*

VSMCs, macrophages, and proliferating and apoptotic cells were identified by immunohistochemistry for  $\alpha$ -smooth muscle cell actin, Mac-2, proliferating cell nuclear antigen (PCNA) and CC-3 as described previously<sup>10</sup>. Fluorescent immunohistochemistry for pAkt was performed using rabbit anti-pAkt antibody (Cell Signalling) diluted 1:25. Fluorescent dual immunohistochemistry for Mac-2 and N-cadherin or FGF-R1 was performed on control atherosclerotic plaques using 5  $\mu$ g/ml rat anti-Mac-2 (Cedar Lane) and 2  $\mu$ g/ml rabbit anti-N-cadherin (Santa Cruz) or rabbit anti-FGF-R1 (Cell Signalling) diluted 1:12.5.

#### *In situ end labelling and actin dual immunohistochemistry*

Apoptotic cells were identified by *in situ* end labelling (ISEL), performed as previously described<sup>12</sup>. This was followed by the smooth muscle  $\alpha$ -actin immunohistochemistry protocol outlined above.

### *Identification of buried fibrous caps*

Serial sections stained for elastin and  $\alpha$ -smooth muscle cell actin were examined for the presence of structures rich in elastin and VSMCs and these were identified as buried fibrous caps, a surrogate marker of previous plaque instability, as previously described<sup>13</sup>.

### **References**

1. Capon DJ, Chamow SM, Mordenti J, Marsters SA, Gregory T, Mitsuya H, Byrn RA, Lucas C, Wurm FM, Groopman JE, Broder S, Smith DH. Designing CD4 immunoadhesins for AIDS therapy. *Nature*. 1989;337:525.
2. Zheng XX, Steele AW, Nickerson PW, Steurer W, Steiger J, Strom TB. Administration of noncytolytic IL-10/Fc in murine models of lipopolysaccharide-induced septic shock and allogeneic islet transplantation. *J Immunol*. 1995;154:5590-5600.
3. Barrou B, Bertry-Coussot L, Morin S, Sainz J, Lucas B, Bitker MO, Debre P, Lemarchand P. Prolonged Islet Allograft Survival by Adenovirus-Mediated Transfer of sICAM-1/Ig Immunoadhesin Gene. *Human Gene Therapy*. 2002;13:1441-1450.
4. Chamow SM, Ashkenazi A. Immunoadhesins: principles and applications. *TIBTECH*. 1996;14:52-60.
5. Chase AJ, Bond M, Crook MF, Newby AC. Role of nuclear factor-kB activation in metalloproteinase-1, -3, and -9 secretion by human macrophages in vitro and rabbit foam cells produced in vivo. *Arterioscler Thromb Vasc Biol*. 2002;22:765-771.
6. Uglow EB, Slater S, Sala-Newby GB, Aguilera-Garcia CM, Angelini GD, Newby AC, George SJ. Dismantling of cadherin-mediated cell-cell contacts modulates smooth muscle proliferation. *Circ Res*. 2003;92:1314-1321.
7. Quasnicka H, Slater SC, Beeching CA, Boehm M, Sala-Newby GB, George SJ. Regulation of smooth muscle cell proliferation by b-catenin/TCF signaling involves modulation of cyclin D1 and p21 expression. *Circ Res*. 2006;99:1329-1337.

8. Shan WS, Yagita Y, Wang Z, Koch A, Svenningsen AF, Gruzglin E, Pedraza L, Colman DR. The minimal essential unit for cadherin-mediated intercellular adhesion comprises extracellular domains 1 and 2. *J Biol Chem*. 2004;279:55914-55923.
9. Reiss K, Maretzky T, Ludwig A, Tousseyn T, de Strooper B, Hartmann D, Saftig P. ADAM10 cleavage of N-cadherin and regulation of cell-cell adhesion and b-catenin nuclear signalling. *EMBO J*. 2005;24:742-752.
10. Johnson JL, Baker AH, Oka K, Chan L, Newby AC, Jackson CL, George SJ. Suppression of atherosclerotic plaque progression and instability by tissue inhibitor of metalloproteinase-2: involvement of macrophage migration and apoptosis. *Circulation*. 2006;113:2435-2444.
11. Johnson J, Jackson C. Atherosclerotic plaque rupture in the apolipoprotein E knockout mouse. *Atherosclerosis*. 2001;154:399-406.
12. George SJ, Johnson J, Angelini G, Newby A, Baker A. Adenovirus-mediated gene transfer of the human TIMP-1 gene inhibits SMC migration and neointima formation in human saphenous vein. *Hum Gene Ther*. 1998;9:867 - 877.
13. Johnson J, Carson K, Williams H, Karanam S, Newby AC, Angelini GD, George SJ, Jackson C. Plaque rupture after short periods of fat feeding in the apolipoprotein E knockout mouse. Model characterization and effects of pravastatin treatment. *Circulation*. 2005;111:1422 - 1430.

## Supplementary Figure Legends

Supplementary Figure I: The pro-survival effect of SNC requires the HAV motif.

Percentage of apoptotic VSMC (CC-3 ICC) 24 hours after FasL treatment with peptide (n=3). \*significant difference from Fc control, §significant difference from HAV.

Supplementary Figure II: Expression of N-cadherin and FGF-R1 in monocytes and macrophages.


Table shows quantitative PCR results for N-cadherin and FGF-R1 in monocytes, macrophages and VSMCs expressed as copy number of mRNA.

Images show representative dual immunohistochemistry for N-cadherin (A, D) or FGF-R1 (B, E) in green and macrophages in red in PBS control atherosclerotic plaque. Arrowheads indicate macrophages expressing N-cadherin or FGF-R. Scale bar in panel A represents 50  $\mu$ m and applies to panels A-C. Scale bar in panel D represents 15  $\mu$ m and applies to panels D-F. Non-immune IgG is shown as negative control (C, F). Nuclei are stained blue with DAPI.

Supplementary Figure III: Schematic diagram of the mechanism of action of SNC


SNC may interact with full length N-cadherin which in turn binds to FGF-R or it may bind directly with FGF-R and full length N-cadherin. These interactions activate PI3-kinase and thereby activate Akt. Active Akt phosphorylates Bad which inhibits the interaction with Bcl-2, released Bcl-2 then provides a survival signal for VSMCs, inhibiting apoptosis.

Supplementary Figure I



## Supplementary Figure II

Copy number of mRNA	N-cadherin	FGF-R1
Monocytes	13.7±13.7 (n=4)	4.2±2.3 (n=3)
Macrophages	1104±449 (n=4)	6.0±4.5 (n=3)
VSMCs	1707±871 (n=6)	168±76 (n=3)





# Supplementary Figure III

